

# UCSF

## UC San Francisco Previously Published Works

### Title

FUS pathology defines the majority of tau- and TDP-43-negative frontotemporal lobar degeneration.

### Permalink

<https://escholarship.org/uc/item/45v4c0cs>

### Journal

Acta neuropathologica, 120(1)

### ISSN

0001-6322

### Authors

Urwin, Hazel  
Josephs, Keith A  
Rohrer, Jonathan D  
et al.

### Publication Date

2010-07-01

### DOI

10.1007/s00401-010-0698-6

Peer reviewed

# FUS pathology defines the majority of tau- and TDP-43-negative frontotemporal lobar degeneration

Hazel Urwin · Keith A. Josephs · Jonathan D. Rohrer · Ian R. Mackenzie · Manuela Neumann · Astrid Authier · Harro Seelaar · John C. Van Swieten · Jeremy M. Brown · Peter Johannsen · Jorgen E. Nielsen · Ida E. Holm · The FReJA Consortium · Dennis W. Dickson · Rosa Rademakers · Neill R. Graff-Radford · Joseph E. Parisi · Ronald C. Petersen · Kimmo J. Hatanpaa · Charles L. White III · Myron F. Weiner · Felix Geser · Vivianna M. Van Deerlin · John Q. Trojanowski · Bruce L. Miller · William W. Seeley · Julie van der Zee · Samir Kumar-Singh · Sebastiaan Engelborghs · Peter P. De Deyn · Christine Van Broeckhoven · Eileen H. Bigio · Han-Xiang Deng · Glenda M. Halliday · Jillian J. Kril · David G. Munoz · David M. Mann · Stuart M. Pickering-Brown · Valerie Doodeman · Gary Adamson · Shabnam Ghazi-Noori · Elizabeth M. C. Fisher · Janice L. Holton · Tamas Revesz · Martin N. Rossor · John Collinge · Simon Mead · Adrian M. Isaacs

Received: 11 January 2010/Revised: 20 April 2010/Accepted: 8 May 2010/Published online: 20 May 2010  
© The Author(s) 2010. This article is published with open access at Springerlink.com

**Abstract** Through an international consortium, we have collected 37 tau- and TAR DNA-binding protein 43 (TDP-43)-negative frontotemporal lobar degeneration (FTLD) cases, and present here the first comprehensive analysis of

these cases in terms of neuropathology, genetics, demographics and clinical data. 92% (34/37) had fused in sarcoma (FUS) protein pathology, indicating that FTLD-FUS is an important FTLD subtype. This FTLD-FUS collection specifically focussed on aFTLD-U cases, one of three recently defined subtypes of FTLD-FUS. The aFTLD-U subtype of

Members of the FReJA consortium are listed in the Appendix.

H. Urwin · A. Authier · G. Adamson · S. Ghazi-Noori · J. Collinge · S. Mead · A. M. Isaacs (✉)  
MRC Prion Unit, UCL Institute of Neurology,  
Queen Square, London WC1N 3BG, UK  
e-mail: a.isaacs@prion.ucl.ac.uk

J. D. Rohrer · M. N. Rossor  
Dementia Research Centre, UCL Institute of Neurology,  
Queen Square, London WC1N 3BG, UK

E. M. C. Fisher · J. Collinge  
Department of Neurodegenerative Disease, UCL Institute  
of Neurology, Queen Square, London WC1N 3BG, UK

J. L. Holton · T. Revesz  
Department of Neuropathology, UCL Institute of Neurology,  
Queen Square, London WC1N 3BG, UK

K. A. Josephs (✉) · R. C. Petersen  
Department of Neurology, Mayo Clinic,  
Rochester, MN, USA  
e-mail: Josephs.Keith@mayo.edu

J. E. Parisi  
Department of Laboratory Medicine and Pathology,  
Mayo Clinic, Rochester, MN, USA

I. R. Mackenzie  
Department of Pathology and Laboratory Medicine,  
Vancouver General Hospital, University of British Columbia,  
Vancouver, BC, Canada

M. Neumann  
Institute of Neuropathology, University Hospital of Zurich,  
Zurich, Switzerland

H. Seelaar · J. C. Van Swieten  
Department of Neurology, Erasmus University Medical Center  
Rotterdam, Rotterdam, The Netherlands

J. M. Brown  
Department of Neurology, Addenbrooke's Hospital,  
Cambridge, UK

P. Johannsen · J. E. Nielsen  
Memory Disorders Research Unit, Department of Neurology,  
Copenhagen University Hospital, Rigshospitalet,  
Copenhagen, Denmark

J. E. Nielsen  
Department of Cellular and Molecular Medicine,  
Section of Neurogenetics, The Panum Institute,  
University of Copenhagen, Copenhagen, Denmark

FTLD-FUS is characterised clinically by behavioural variant frontotemporal dementia (bvFTD) and has a particularly young age of onset with a mean of 41 years. Further, this subtype had a high prevalence of psychotic symptoms (36% of cases) and low prevalence of motor symptoms (3% of cases). We did not find *FUS* mutations in any aFTLD-U case. To date, the only subtype of cases reported to have ubiquitin-positive but tau-, TDP-43- and FUS-negative pathology, termed FTLD-UPS, is the result of charged multivesicular body protein 2B gene (*CHMP2B*) mutation. We identified three FTLD-UPS cases, which are negative for *CHMP2B* mutation, suggesting that the full complement of FTLD pathologies is yet to be elucidated.

**Keywords** FTLD · FUS · FTLD-UPS · Frontotemporal · FTD

### Abbreviations

bvFTD Behavioural variant FTD  
CHMP2B Charged multivesicular body protein 2B

FTD	Frontotemporal dementia
FTD-3	FTD linked to chromosome 3
FTLD	Frontotemporal lobar degeneration
FTLD-IF	Frontotemporal lobar degeneration with intermediate filament positive inclusions
FTLD-ni	Frontotemporal lobar degeneration with no inclusions
FTLD-TDP	Frontotemporal lobar degeneration with TDP-43 positive inclusions
FTLD-UPS	Frontotemporal dementia with ubiquitin positive, TDP-43 negative inclusions
GRN	Progranulin
MAPT	Microtubule associated protein tau
NCI	Neuronal cytoplasmic inclusion
NII	Neuronal intranuclear inclusion
PPA	Primary progressive aphasia
TDP-43	Transactive response binding protein of 43 kDa
VCP	Valosin containing protein

I. E. Holm  
Department of Pathology, Randers Hospital,  
Randers and Laboratory for Experimental Neuropathology,  
Danish Neuroscience Center, Aarhus University Hospital,  
Aarhus, Denmark

D. W. Dickson · R. Rademakers · N. R. Graff-Radford  
Department of Neuroscience, Mayo Clinic, Jacksonville,  
FL, USA

K. J. Hatanpaa · C. L. White III  
Department of Pathology, University of Texas Southwestern  
Medical School, Dallas, TX, USA

M. F. Weiner  
Departments of Psychiatry and Neurology, University of Texas  
Southwestern Medical School, Dallas, TX, USA

F. Geser · V. M. Van Deerlin · J. Q. Trojanowski  
Center for Neurodegenerative Disease Research, Department  
of Pathology and Laboratory Medicine, University of  
Pennsylvania School of Medicine, Philadelphia, USA

B. L. Miller · W. W. Seeley  
UCSF Department of Neurology, Memory and Aging Center,  
San Francisco, CA, USA

J. van der Zee · S. Kumar-Singh · C. Van Broeckhoven  
VIB Department of Molecular Genetics, Neurodegenerative  
Brain Diseases Group, Antwerpen, Belgium

J. van der Zee · S. Kumar-Singh · S. Engelborghs ·  
P. P. De Deyn · C. Van Broeckhoven  
Institute Born-Bunge, University of Antwerp,  
Antwerpen, Belgium

S. Engelborghs · P. P. De Deyn  
Department of Neurology, Memory Clinic, ZNA Middelheim,  
Antwerpen, Belgium

E. H. Bigio  
Department of Pathology, Feinberg School of Medicine,  
Northwestern University, Chicago, IL, USA

H.-X. Deng  
Department of Neurology and Clinical Neurosciences,  
Feinberg School of Medicine, Northwestern University,  
Chicago, IL, USA

G. M. Halliday  
Prince of Wales Medical Research Institute,  
University of New South Wales,  
Sydney, NSW, Australia

J. J. Kril  
Disciplines of Medicine and Pathology, University of Sydney,  
Sydney, NSW, Australia

D. G. Munoz  
Department of Pathology, Saint Michael's Hospital,  
University of Toronto, Toronto, ON, Canada

D. M. Mann  
Clinical Neuroscience Research Group, School of Translational  
Medicine, Greater Manchester Neurosciences Centre,  
University of Manchester, Salford, UK

S. M. Pickering-Brown  
Clinical Neurosciences Research Group, Faculty of Human  
and Medical Sciences, University of Manchester,  
Oxford Road, Manchester M13 9PT, UK

V. Doodeman  
Department of Clinical Genetics, VU University Medical Center,  
Amsterdam, The Netherlands

## Introduction

Frontotemporal lobar degeneration (FTLD) describes a group of diseases characterised by bilateral, often asymmetric, atrophy of the frontal and anterior temporal lobes. Frontotemporal dementia (FTD), also termed behavioural variant FTD (bvFTD), is the most common clinical manifestation, but FTLD can also cause the language disorders progressive non-fluent aphasia (PNFA) and semantic dementia (SD), collectively known as primary progressive aphasia (PPA). The clinical presentation is related to the distribution of pathology in the frontal and temporal lobes. Despite gross pathological similarities, FTLD comprises a clinically, genetically, and neuropathologically heterogeneous collection of disorders.

A family history of a similar disease has been reported in 25–40% of FTLD cases [6, 44, 50], although this estimate is much lower in some populations [17]. FTLD-causing mutations have been found in the microtubule-associated protein tau (*MAPT*) [16], progranulin (*GRN*) [1, 7] and charged multivesicular body protein 2B (*CHMP2B*) genes [49]. Mutations in valosin-containing protein (*VCP*) cause inclusion body myopathy and Paget disease of bone with frontotemporal dementia, in which 20–100% of patients can develop FTD [10, 53, 56]. *GRN* mutations account for approximately 10% of FTLD cases [2, 7–9] and *MAPT* up to 10% depending on the population studied [15, 41, 48]; while mutations in *CHMP2B* are a much rarer cause of FTLD, having been reported in one Danish family termed frontotemporal dementia linked to chromosome 3 (FTD-3) [25, 49, 52] and one unrelated Belgian familial FTLD patient [54].

Up to 40% of FTLD cases, including all *MAPT* mutation cases, have tau pathology [48] (FTLD-tau); while over 50% have neuronal inclusions, which are immunoreactive for ubiquitin in the absence of abnormal tau,  $\alpha$ -synuclein, or amyloid deposition [20]. In the majority of tau-negative cases, the major ubiquitinated protein is the TAR DNA-binding protein 43 (TDP-43) [11, 39], and these are designated FTLD-TDP [30]. FTLD-TDP can be further classified into neuropathological subtypes depending on the distribution, morphology and precise cellular location of inclusions [28, 45].

Recent studies report that 6–20% of tau-negative FTLD cases have ubiquitinated inclusions which are TDP-43 negative [12, 21, 29, 42]. Ubiquitin positive, tau and TDP-43 negative cases include those with neuronal intermediate filament- and  $\alpha$ -internexin-positive inclusions [5, 19], termed NIFID, and basophilic inclusion body disease (BIBD) [35, 58]. Distinct from these are a subset of FTLD cases termed atypical FTLD-U (aFTLD-U), which do not have intermediate filament positive or basophilic inclusions, but are characterised by unusual curved or twisted

neuronal intranuclear ubiquitin positive inclusions, as well as cytoplasmic inclusions [29, 42]. The recent finding of FUS (fused in sarcoma) protein pathology in aFTLD-U cases [37] and subsequently in NIFID and BIBD cases [34, 38] has led to further reorganisation of FTLD classification [31]: aFTLD-U, NIFID and BIBD are now considered three distinct subtypes of FTLD-FUS, the collective term for FTLD cases with FUS pathology. Cases with ubiquitinated inclusions that are negative for tau, TDP-43 and FUS are termed FTLD-UPS [30]; this includes FTLD cases caused by *CHMP2B* mutations [13, 14].

FUS pathology was first described in familial ALS cases following the discovery of causative mutations in the *FUS* gene [23, 55]. FUS is a multifunctional DNA- and RNA-binding protein, sharing functional homology with TDP-43 [24]. FUS pathology has also been found in polyglutamine expansion diseases including Huntington's disease, suggesting a broad spectrum of FUS proteinopathies [34, 38, 57]. However, FUS pathology was not found in FTD-3, a major subtype of FTLD-UPS [14].

A minority of cases do not have tau, TDP-43, FUS or ubiquitinated inclusions and these have been termed FTLD with no inclusions (FTLD-ni, previously dementia lacking distinctive histopathology, or DLDH) [30]. Increasingly sensitive immunostaining protocols have led to most FTLD-ni cases being reclassified [20, 26, 32]; however, a small group remain negative for characterised inclusions despite the use of updated ubiquitin immunohistochemistry.

Here, we report, through a major international collaboration, the first comprehensive collection of the aFTLD-U subtype of FTLD-FUS and describe their clinical and demographic features.

## Materials and methods

### Cases

A total of 37 cases with ubiquitin-positive but tau- and TDP-43-negative inclusions were evaluated from 12 academic research centres. In each centre, clinical information was acquired from case notes by a neurologist with expertise in neurodegenerative diseases. Standard neurological and cognitive assessments were used to ascertain the presence of behavioural, cognitive and motor abnormalities. All centres followed consensus clinical criteria for FTLD diagnosis [33, 36], and were asked to supply presenting symptoms for each patient and the final clinical diagnosis that had been rendered by the evaluating neurologist. These data were collated and then reviewed by an expert in FTLD (K.A.J.) to ensure consistency in diagnosis among the different centres. BIBD and NIFID cases were specifically excluded in order to focus on a specific subtype

of FTL-D-FUS. Twenty-three cases for which DNA was available (21 FTL-D-FUS and 2 FTL-D-UPS) were screened for *CHMP2B* mutations. Fifteen FTL-D-FUS cases were screened for *FUS* mutations and a further six have been reported previously as *FUS* mutation negative [37]. Patients were considered to have a family history if a first- or second-degree relative had dementia. This study was approved by the UCL Institute of Neurology and National Hospital for Neurology and Neurosurgery Local Research Ethics Committee.

### Histopathology

All cases were stained for TDP-43, ubiquitin and  $\alpha$ -internexin or neurofilament proteins according to local protocols at each centre [2, 12, 21, 27, 29, 39, 40, 42, 47, 53]. Experienced neuropathologists in each centre analysed the sections to give the neuropathological diagnosis. FUS staining was subsequently performed on frontal cortex and hippocampus for all cases. As previously described, FUS staining is sensitive to length of fixation [37]. Therefore, at least two antibodies (Sigma HPA008784 and Bethyl Laboratories A300-302A) were used for all cases examined and antibody concentrations optimised for each case by ensuring physiological FUS staining was visible, as recommended in the original description of FTL-D-FUS [37]. To further ensure inter-centre reliability 26 cases, including the three remaining FTL-D-UPS cases, were stained in one of two laboratories which first reported FTL-D-FUS (IRM or MN).

### Genetic analysis

Genomic DNA was extracted from brain tissue using standard phenol–chloroform extraction, or DNA extracted from brain or blood was directly provided by the contributor. The entire open reading frame and exon–intron boundaries of *CHMP2B* and *FUS* were sequenced, except for three cases in which *FUS* exons 14 and 15 only were sequenced. Primers and conditions for *CHMP2B* sequencing have been described previously [49]; conditions for *FUS* sequencing are available on request. Sequences were analysed using the SeqScape software Version 2.5 (Applied Biosystems, Foster City, CA, USA).

## Results

### Neuropathology

A total of 37 ubiquitin-positive, tau- and TDP-43-negative cases were examined. 92% (34/37) of cases were characterised by neuronal cytoplasmic and intranuclear inclusions (NIIs) which were immunoreactive for FUS, corresponding to the aFTLD-U subtype of FTL-D-FUS. The NIIs were as

previously described, having an unusual morphology, which can be long and straight, curved or twisted [29, 42]. Striatal degeneration was present in 100% (27/27) of aFTLD-U cases and hippocampal sclerosis was present in 97% (29/30) of cases. 8% (3/37) were characterised by ubiquitin-positive neuronal cytoplasmic inclusions (NCIs) which were FUS-negative, i.e. FTL-D-UPS.

### Genetics

Sequencing of the *CHMP2B* gene in all cases for which DNA was available revealed no further coding mutations. Two synonymous coding variants were identified in three FTL-D-FUS cases: c.27C>T, p.Thr9Thr and c.372A>C, p.Thr124Thr. These two variants were each observed in the heterozygous state but always together, suggesting they could be in linkage disequilibrium. None of these variants has been previously reported, but as they do not lead to an amino acid change, they are unlikely to be pathogenic. Therefore, *CHMP2B* mutation-positive cases comprise a distinct genetic subgroup of FTL-D-UPS. No mutations were identified in *FUS* in any case.

### Demographics

#### Onset

Age at onset data was available for 33 aFTLD-U cases. The mean age of onset was 41.2 years (standard deviation 9.3 years). The three FTL-D-UPS cases had a mean age of onset of 50.3 years (standard deviation 4 years).

#### Disease duration

Disease duration data were available for 33 aFTLD-U cases. Mean disease duration was 7.7 years (standard deviation 3.2 years). Disease duration data were available for only two of the three FTL-D-UPS cases and were 4 and 11 years, respectively.

#### Sex

Of the aFTLD-U cases, 47% (16/34) were female and all three FTL-D-UPS cases were female.

#### Family history

One aFTLD-U case (3%, 1/34) had a family history and 2/3 of the FTL-D-UPS cases had positive family histories.

### Clinical features

The clinical features of several aFTLD-U cases have previously been reported [21, 29, 42, 46] and analysed together,

**Table 1** Clinical symptoms in aFTLD-U cases

ID	Sex	Diagnosis	Behavioural/personality					Psychiatric		Cognitive	Motor	
			Apathy	Disinh	Abnormal eating	Compulsions	Inappropriate sexual	Delusions	Hallucinations	Aphasia	Parkinsonism	MND
UBC1	F	bvFTD	1	1	1	1	0	0	0	0	0	0
UBC2	F	bvFTD	1	0	0	0	1	1	0	0	0	0
UBC3	F	bvFTD	1	1	1	0	1	0	0	0	0	0
UBC4	F	bvFTD	1	1	1	0	0	0	0	0	0	0
UBC5	M	bvFTD	1	1	1	0	1	1	0	0	0	0
UBC6	F	bvFTD	1	1	0	0	0	0	1	0	0	0
MUC1	M	Dem	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
MUC2	F	bvFTD	1	1	0	0	NA	NA	NA	1	0	0
MUC3	F	Dem	NA	1	NA	NA	NA	NA	NA	NA	NA	NA
MUC4	M	bvFTD	1	1	1	1	NA	NA	NA	1	0	0
MUC5	F	bvFTD	0	1	1	0	NA	NA	NA	1	0	0
MUC6	M	bvFTD	1	1	NA	NA	NA	NA	NA	NA	NA	NA
MUC7	F	bvFTD	1	1	1	0	0	0	0	0	0	0
MUC8	M	bvFTD	NA	1	NA	NA	NA	NA	NA	NA	NA	NA
MUC9	M	bvFTD	0	1	1	1	NA	NA	NA	0	0	0 <sup>a</sup>
UR1	M	bvFTD	1	1	1	1	1	0	1	0	0	0
UR2	F	bvFTD	1	1	1	0	1	0	0	0	0	0
UR3	F	bvFTD	1	NA	1	1	0	1	0	0	0	0
UR4	F	bvFTD	1	1	0	1	0	0	0	0	0	0
UT1	M	bvFTD	NA	1	NA	NA	NA	1	0	0	0	0
UTSW1	M	bvFTD	1	1	1	1	1	0	0	1	0	0
NWU1	M	bvFTD	0	1	1	1	1	0	0	0	0	0
NWU2	F	bvFTD	1	1	1	1	1	0	0	1	0	0
UP1	F	bvFTD	1	1	1	1	0	NA	NA	0	0	0
UP2	M	bvFTD	1	0	1	1	0	NA	NA	0	0	0
UP3	M	bvFTD	1	1	NA	1	0	NA	NA	0	0	0
UNSW1	F	bvFTD	NA	1	NA	1	NA	0	0	1	1	0
UNSW2	M	bvFTD	NA	1	1	0	0	0	0	1	0	0
MC1	M	bvFTD	0	1	1	1	1	0	1	0	0	0
MC2	F	bvFTD	0	1	NA	1	1	NA	NA	0	0	0
MC3	M	bvFTD	0	1	1	1	1	0	0	0	0	0
MC4	M	bvFTD	1	0	0	1	0	0	0	0	0	0
MC5	M	bvFTD	0	1	1	0	1	0	0	0	0	0
UCL1	M	bvFTD	NA	1	1	NA	NA	0	1	NA	0	0
% with symptom			74	91	81	61	52	18	18	24	3	0

Cases of aFTLD-U were scored for presence of the symptoms indicated

0 absent, 1 present, NA information not available, *bvFTD* behavioural variant FTD, *Dem* unspecified dementia, *Disinh* disinhibition

<sup>a</sup> One patient exhibited tongue fasciculation, dysphagia and muscle hypotonia but there was insufficient evidence to diagnose MND

these cases form a clinically homogeneous group. 94% (32/34) of aFTLD-U patients received an initial diagnosis of bvFTD (Table 1). The other two patients, for whom there was little clinical information available, were described as having ‘unspecified dementia’. Patients were typically described as showing a decline in social conduct, self-neglect and disinhibition, with aggression as a common

feature. Inappropriate sexual behaviour was widely reported, in 52% of cases (12/23) (Table 1). Dietary changes, most commonly overeating, were observed in 81% of cases (21/26) with hyperorality a feature of several.

Strikingly, psychotic symptoms, either hallucinations or delusions, were observed in 36% of aFTLD-U patients (8/22), and not comorbid in any case. Motor symptoms

were infrequent and when present, they were typically limited to mild rigidity or intermittent hyperkinesias. Only one patient (1/30, 3%) showed classical parkinsonism.

The three FTLD-UPS cases all had an initial diagnosis of bvFTD. Very little clinical information was available, but apathy was common to two of these cases as was parkinsonism.

## Discussion

This study describes the first comprehensive collection of the aFTLD-U subtype of FTLD-FUS. We show that aFTLD-U has a distinct clinical phenotype characterised by the behavioural variant FTD syndrome, a young onset of disease, a high prevalence of psychotic symptoms and a low prevalence of motor symptoms. FUS pathology accounted for 92% of cases with ubiquitin-positive, tau- and TDP-43-negative FTLD showing that it defines the vast majority of such cases. However, we identified three FTLD-UPS cases without CHMP2B mutation showing that while such cases are rare, the FTLD-UPS classification has not been completely subsumed by CHMP2B mutation cases or FTLD-FUS.

Of all of the FTLD subtypes, bvFTD is pathologically the most heterogeneous making it difficult to predict the underlying pathology from clinical features in life. This cohort similarly has an overlapping range of symptoms with other sporadic and genetic causes of bvFTD. However, psychotic symptoms are rarely described in other cases of bvFTD [22], and the presence of delusions and hallucinations in a substantial proportion of this cohort may help to distinguish them from other patients. The very low frequency of motor symptoms may also help define this group of patients.

The mean age of onset of 41.2 years for the aFTLD-U cases is much lower than that of a previously reported sporadic FTLD cohort (60.5 years) [4], and therefore, may help distinguish them from other sporadic FTLD cases. 15/34 aFTLD-U cases came from two contributing centres, therefore, it is possible that the results have been biased by these cases, which were previously reported to have a low age of onset [29, 42]. However, the age of onset of these cases is not significantly different from the other 19 cases in the study (*t* test, ns). Furthermore, psychotic symptoms are evenly distributed between cases from these two centres (3/7, 42%) and cases from other centres (5/15, 33%) (Fisher's exact test, ns).

Only one aFTLD-U case (3%, 1/34) had a positive family history for FTD: this patient was one of nine siblings, and had one brother who was diagnosed clinically with 'Pick's disease' at 61 years of age and died at 64 years of age without autopsy. Another brother was reported to show 'cognitive changes' at the age of 64 years.

The patient was negative for *FUS* mutation, suggesting that there may be other genetic causes of FTLD-FUS. While aFTLD-U cases have a distinctive and consistent clinicopathological phenotype, FTLD appears to occur sporadically in the majority of cases. Of the three FTLD-UPS cases, two had a family history, and one had at least two relatives affected with the same disease [43]. These cases were negative for CHMP2B mutations, which suggests other genetic causes of FTLD-UPS and the possibility of at least one other unidentified ubiquitinated protein associated with FTLD. No gender bias was found in any of the groups studied. A female preponderance was previously reported in an analysis of a subset of aFTLD-U cases [29]; however, this was not confirmed in wider analysis.

It has recently been shown that two rare subtypes of FTLD, NIFID and BIBD have extensive FUS pathology [34, 38]. We did not include NIFID and BIBD cases in this study as they have been described to have distinct clinical features to aFTLD-U cases, particularly prominent motor symptoms [3, 18, 19, 34, 51, 58]. In order to allow comparison of FTLD-FUS to other FTLD cases, we focussed on one subtype of FTLD-FUS (aFTLD-U), as it is possible that FTLD-FUS comprises a clinically heterogeneous group. However, now that NIFID, BIBD and aFTLD-U have been shown to have a common underlying pathology, it will be interesting to directly compare these different FUS entities in future studies. FUS pathology is also found in polyglutamine expansion diseases including Huntington's disease [57], showing that it is not limited to the FTLD-MND spectrum of diseases. This is similar to tau and TDP-43 pathology, which are pathological hallmarks of different FTLD subtypes but are also observed in a wide range of other neurodegenerative diseases. Within the clinical syndrome of FTLD, FUS pathology will be an extremely important tool for the classification of cases which are tau- and TDP-43- negative, and wider analysis of FTLD-FUS cases may reveal further subtypes.

In conclusion, the vast majority of cases with tau- and TDP-43 negative inclusions have FUS pathology, suggesting that FTLD-FUS is an important FTLD subtype. The opportunity to analyse a large series of aFTLD-U cases has only been possible through a large multi-centre collaboration, with all centres contributing a relatively small number of cases. This collaborative effort has provided new insights into the genetic and clinicopathological spectrum of FTLD-FUS and FTLD-UPS.

**Conflict of interest statement** The authors report no conflicts of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



## Appendix: The Frontotemporal Dementia Research in Jutland Association (FReJA)

Anders Gade, Tove Thusgaard, Memory Disorders Research Group, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark. Susanne Gydesen, Psychiatric Centre Ballerup, Copenhagen University Hospital, Ballerup, Denmark. Elisabet Englund, Department of Pathology, Lund University Hospital, Lund, Sweden. This work was undertaken at UCLH/UCL, and was supported by the Department of Health's NIHR Biomedical Research Centres; the Medical Research Council, UK; the National Institute on Aging, National Institutes of Health [grant numbers P30 AG012300 to UTSW, P30 AG13854 to NWU, P50 AG16574 to Mayo Clinic, AG10124 and AG17546 to U. Penn.]; the National Center for Research Resources, National Institutes of Health [grant number UL1RR024982]; the National Institutes of Health [grant number NIH NS65782-01 J]; the Fund for Scientific Research Flanders (FWO-V); the IAP P6/43 network of the Belgian Science Policy Office (BELSPO); the Winspear Family Center for Research on the Neuropathology of Alzheimer Disease and the McCune Foundation.

## References

- Baker M, Mackenzie IR, Pickering-Brown SM, Gass J, Rademakers R, Lindholm C, Snowden J, Adamson J, Sadovnick AD, Rollinson S, Cannon A, Dwosh E, Neary D, Melquist S, Richardson A, Dickson D, Berger Z, Eriksen J, Robinson T, Zehr C, Dickey CA, Crook R, McGowan E, Mann D, Boeve B, Feldman H, Hutton M (2006) Mutations in progranulin cause tau-negative frontotemporal dementia linked to chromosome 17. *Nature* 442:916–919
- Beck J, Rohrer JD, Campbell T, Isaacs A, Morrison KE, Goodall EF, Warrington EK, Stevens J, Revesz T, Holton J, Al Sarraj S, King A, Scahill R, Warren JD, Fox NC, Rossor MN, Collinge J, Mead S (2008) A distinct clinical, neuropsychological and radiological phenotype is associated with progranulin gene mutations in a large UK series. *Brain* 131:706–720
- Cairns NJ, Grossman M, Arnold SE, Burn DJ, Jaros E, Perry RH, Duyckaerts C, Stankoff B, Pillon B, Skullerud K, Cruz-Sanchez FF, Bigio EH, Mackenzie IR, Gearing M, Juncos JL, Glass JD, Yokoo H, Nakazato Y, Mosaheb S, Thorpe JR, Uryu K, Lee VM, Trojanowski JQ (2004) Clinical and neuropathologic variation in neuronal intermediate filament inclusion disease. *Neurology* 63:1376–1384
- Cairns NJ, Neumann M, Bigio EH, Holm IE, Troost D, Hatanpaa KJ, Foong C, White CL III, Schneider JA, Kretschmar HA, Carter D, Taylor-Reinwald L, Paulsmeier K, Strider J, Gitcho M, Goate AM, Morris JC, Mishra M, Kwong LK, Stieber A, Xu Y, Forman MS, Trojanowski JQ, Lee VM, Mackenzie IR (2007) TDP-43 in familial and sporadic frontotemporal lobar degeneration with ubiquitin inclusions. *Am J Pathol* 171:227–240
- Cairns NJ, Uryu K, Bigio EH, Mackenzie IRA, Gearing M, Duyckaerts C, Yokoo H, Nakazato Y, Jaros E, Perry RH, Arnold SE, Lee VM, Trojanowski JQ (2004) alpha-Internexin aggregates are abundant in neuronal intermediate filament inclusion disease (NIFID) but rare in other neurodegenerative diseases. *Acta Neuropathol* 108:213–223
- Chow TW, Miller BL, Hayashi VN, Geschwind DH (1999) Inheritance of frontotemporal dementia. *Arch Neurol* 56:817–822
- Cruts M, Gijselinck I, van der Zee J, Engelborghs S, Wils H, Pirici D, Rademakers R, Vandenberghe R, Dermaut B, Martin JJ, van Duijn C, Peeters K, Sciot R, Santens P, De Pooter T, Mattheijssens M, Van den Broeck M, Cuijt I, Vennekens K, De Deyn PP, Kumar-Singh S, Van Broeckhoven C (2006) Null mutations in progranulin cause ubiquitin-positive frontotemporal dementia linked to chromosome 17q21. *Nature* 442:920–924
- Cruts M, Van Broeckhoven C (2008) Loss of progranulin function in frontotemporal lobar degeneration. *Trends Genet* 24:186–194
- Gass J, Cannon A, Mackenzie IR, Boeve B, Baker M, Adamson J, Crook R, Melquist S, Kuntz K, Petersen R, Josephs K, Pickering-Brown SM, Graff-Radford N, Uitti R, Dickson D, Wszolek Z, Gonzalez J, Beach TG, Bigio E, Johnson N, Weintraub S, Mesulam M, White CL III, Woodruff B, Caselli R, Hsiung GY, Feldman H, Knopman D, Hutton M, Rademakers R (2006) Mutations in progranulin are a major cause of ubiquitin-positive frontotemporal lobar degeneration. *Hum Mol Genet* 15:2988–3001
- Guyant-Marechal L, Laquerriere A, Duyckaerts C, Dumanchin C, Bou J, Dugny F, Le Ber I, Frebourg T, Hannequin D, Campion D (2006) Valosin-containing protein gene mutations: clinical and neuropathologic features. *Neurology* 67:644–651
- Hasegawa M, Arai T, Nonaka T, Kametani F, Yoshida M, Hashizume Y, Beach TG, Buratti E, Baralle F, Morita M, Nakano I, Oda T, Tsuchiya K, Akiyama H (2008) Phosphorylated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Ann Neurol* 64:60–70
- Hatanpaa KJ, Bigio EH, Cairns NJ, Womack KB, Weintraub S, Morris JC, Foong C, Xiao G, Hladik C, Manton TY, White CL III (2008) TAR DNA-binding protein 43 immunohistochemistry reveals extensive neuritic pathology in FTL-DU: a midwest-southwest consortium for FTL-DU study. *J Neuropathol Exp Neurol* 67:271–279
- Holm IE, Englund E, Mackenzie IR, Johannsen P, Isaacs AM (2007) A reassessment of the neuropathology of frontotemporal dementia linked to chromosome 3. *J Neuropathol Exp Neurol* 66:884–891
- Holm IE, Isaacs AM, Mackenzie IRA (2009) Absence of FUS-immunoreactive pathology in frontotemporal dementia linked to chromosome 3 (FTD-3) caused by mutation in the CHMP2B gene. *Acta Neuropathol* 118:719–720
- Houlden H, Baker M, Adamson J, Grover A, Waring S, Dickson D, Lynch T, Boeve B, Petersen RC, Pickering-Brown S, Owen F, Neary D, Craufurd D, Snowden J, Mann D, Hutton M (1999) Frequency of tau mutations in three series of non-Alzheimer's degenerative dementia. *Ann Neurol* 46:243–248
- Hutton M, Lendon CL, Rizzo P, Baker M, Froelich S, Houlden H, Pickering-Brown S, Chakraverty S, Isaacs A, Grover A, Hackett J, Adamson J, Lincoln S, Dickson D, Davies P, Petersen RC, Stevens M, de Graaff E, Wauters E, van Baren J, Hillebrand M, Joosse M, Kwon JM, Nowotny P, Che LK, Norton J, Morris JC, Reed LA, Trojanowski J, Basun H, Lannfelt L, Neystat M, Fahn S, Dark F, Tannenberg T, Dodd PR, Hayward N, Kwok JB, Schofield PR, Andreadis A, Snowden J, Craufurd D, Neary D, Owen F, Oostra BA, Hardy J, Goate A, van Swieten J, Mann D, Lynch T, Heutink P (1998) Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 393:702–705



17. Ikeda K (2000) Neuropathologic discrepancy between Japanese Pick's disease without Pick bodies and frontal lobe degeneration type of frontotemporal dementia proposed by Lund and Manchester Group. *Neuropathology* 20:76–82
18. Ishihara K, Araki S, Ihori N, Shiota J, Kawamura M, Nakano I (2006) An autopsy case of frontotemporal dementia with severe dysarthria and motor neuron disease showing numerous basophilic inclusions. *Neuropathology* 26:447–454
19. Josephs KA, Holton JL, Rossor MN, Braendgaard H, Ozawa T, Fox NC, Petersen RC, Pearl GS, Ganguly M, Rosa P, Laursen H, Parisi JE, Waldemar G, Quinn NP, Dickson DW, Revesz T (2003) Neurofilament inclusion body disease: a new proteinopathy? *Brain* 126:2291–2303
20. Josephs KA, Holton JL, Rossor MN, Godbolt AK, Ozawa T, Strand K, Khan N, Al-Sarraj S, Revesz T (2004) Frontotemporal lobar degeneration and ubiquitin immunohistochemistry. *Neuropathol Appl Neurobiol* 30:369–373
21. Josephs KA, Lin WL, Ahmed Z, Stroh DA, Graff-Radford NR, Dickson DW (2008) Frontotemporal lobar degeneration with ubiquitin-positive, but TDP-43-negative inclusions. *Acta Neuropathol* 116:159–167
22. Knibb JA, Kipps CM, Hodges JR (2006) Frontotemporal dementia. *Curr Opin Neurol* 19:565–571
23. Kwiakowski TJ Jr, Bosco DA, Leclerc AL, Tamrazian E, Vanderburg CR, Russ C, Davis A, Gilchrist J, Kasarskis EJ, Munsat T, Valdmanis P, Rouleau GA, Hosler BA, Cortelli P, de Jong PJ, Yoshinaga Y, Haines JL, Pericak-Vance MA, Yan J, Ticozzi N, Siddique T, Kenna-Yasek D, Sapp PC, Horvitz HR, Landers JE, Brown RH Jr (2009) Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323:1205–1208
24. Lagier-Tourenne C, Cleveland DW (2009) Rethinking ALS: the FUS about TDP-43. *Cell* 136:1001–1004
25. Lindquist SG, Braedgaard H, Svenstrup K, Isaacs AM, Nielsen JE (2008) Frontotemporal dementia linked to chromosome 3 (FTD-3)—current concepts and the detection of a previously unknown branch of the Danish FTD-3 family. *Eur J Neurol* 15:667–670
26. Lipton AM, White CL III, Bigio EH (2004) Frontotemporal lobar degeneration with motor neuron disease-type inclusions predominates in 76 cases of frontotemporal degeneration. *Acta Neuropathol* 108:379–385
27. Luty AA, Kwok JB, Thompson EM, Blumbergs P, Brooks WS, Loy CT, Dobson-Stone C, Panegyres PK, Hecker J, Nicholson GA, Halliday GM, Schofield PR (2008) Pedigree with frontotemporal lobar degeneration—motor neuron disease and Tar DNA binding protein-43 positive neuropathology: genetic linkage to chromosome 9. *BMC Neurol* 8:32
28. Mackenzie IR, Baborie A, Pickering-Brown S, Du PD, Jaros E, Perry RH, Neary D, Snowden JS, Mann DM (2006) Heterogeneity of ubiquitin pathology in frontotemporal lobar degeneration: classification and relation to clinical phenotype. *Acta Neuropathol* 112:539–549
29. Mackenzie IR, Foti D, Woulfe J, Hurwitz TA (2008) Atypical frontotemporal lobar degeneration with ubiquitin-positive, TDP-43-negative neuronal inclusions. *Brain* 131:1282–1293
30. Mackenzie IR, Neumann M, Bigio EH, Cairns NJ, Alafuzoff I, Kril J, Kovacs GG, Ghetti B, Halliday G, Holm IE, Ince PG, Kamphorst W, Revesz T, Rozemuller AJ, Kumar-Singh S, Akiyama H, Baborie A, Spina S, Dickson DW, Trojanowski JQ, Mann DM (2009) Nomenclature for neuropathologic subtypes of frontotemporal lobar degeneration: consensus recommendations. *Acta Neuropathol* 117:15–18
31. Mackenzie IR, Neumann M, Bigio EH, Cairns NJ, Alafuzoff I, Kril J, Kovacs GG, Ghetti B, Halliday G, Holm IE, Ince PG, Kamphorst W, Revesz T, Rozemuller AJ, Kumar-Singh S, Akiyama H, Baborie A, Spina S, Dickson DW, Trojanowski JQ, Mann DM (2010) Nomenclature and nosology for neuropathologic subtypes of frontotemporal lobar degeneration: an update. *Acta Neuropathol* 119:1–4
32. Mackenzie IR, Shi J, Shaw CL, Duplessis D, Neary D, Snowden JS, Mann DM (2006) Dementia lacking distinctive histology (DLDH) revisited. *Acta Neuropathol* 112:551–559
33. McKhann GM, Albert MS, Grossman M, Miller B, Dickson D, Trojanowski JQ (2001) Clinical and pathological diagnosis of frontotemporal dementia: report of the Work Group on Frontotemporal Dementia and Pick's Disease. *Arch Neurol* 58:1803–1809
34. Munoz DG, Neumann M, Kusaka H, Yokota O, Ishihara K, Terada S, Kuroda S, Mackenzie IR (2009) FUS pathology in basophilic inclusion body disease. *Acta Neuropathol* 118:617–627
35. Munoz-Garcia D, Ludwin SK (1984) Classic and generalized variants of Pick's disease: a clinicopathological, ultrastructural, and immunocytochemical comparative study. *Ann Neurol* 16:467–480
36. Neary D, Snowden JS, Gustafson L, Passant U, Stuss D, Black S, Freedman M, Kertesz A, Robert PH, Albert M, Boone K, Miller BL, Cummings J, Benson DF (1998) Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 51:1546–1554
37. Neumann M, Rademakers R, Roeber S, Baker M, Kretschmar HA, Mackenzie IR (2009) A new subtype of frontotemporal lobar degeneration with FUS pathology. *Brain* 132:2922–2931
38. Neumann M, Roeber S, Kretschmar HA, Rademakers R, Baker M, Mackenzie IR (2009) Abundant FUS-immunoreactive pathology in neuronal intermediate filament inclusion disease. *Acta Neuropathol* 118:605–616
39. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, Bruce J, Schuck T, Grossman M, Clark CM, McCluskey LF, Miller BL, Masliah E, Mackenzie IR, Feldman H, Feiden W, Kretschmar HA, Trojanowski JQ, Lee VM (2006) Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* 314:130–133
40. Pirici D, Vandenberghe R, Rademakers R, Dermaut B, Cruts M, Vennekens K, Cuijt I, Lubke U, Ceuterick C, Martin JJ, Van Broeckhoven C, Kumar-Singh S (2006) Characterization of ubiquitinated intraneuronal inclusions in a novel Belgian frontotemporal lobar degeneration family. *J Neuropathol Exp Neurol* 65:289–301
41. Rademakers R, Cruts M, Van Broeckhoven C (2004) The role of tau (MAPT) in frontotemporal dementia and related tauopathies. *Hum Mutat* 24:277–295
42. Roeber S, Mackenzie IR, Kretschmar HA, Neumann M (2008) TDP-43-negative FTL-D-U is a significant new clinico-pathological subtype of FTL-D. *Acta Neuropathol* 116:147–157
43. Rohrer JD, Guerreiro R, Vandrovicova J, Uphill J, Reiman D, Beck J, Isaacs AM, Authier A, Ferrari R, Fox NC, Mackenzie IR, Warren JD, de Silva R, Holton J, Revesz T, Hardy J, Mead S, Rossor MN (2009) The heritability and genetics of frontotemporal lobar degeneration. *Neurology* 73:1451–1456
44. Rosso SM, Donker KL, Baks T, Joosse M, de Koning I, Pijnenburg Y, de Jong D, Dooijes D, Kamphorst W, Ravid R, Niermeijer MF, Verheij F, Kremer HP, Scheltens P, van Duijn CM, Heutink P, van Swieten JC (2003) Frontotemporal dementia in The Netherlands: patient characteristics and prevalence estimates from a population-based study. *Brain* 126:2016–2022
45. Sampathu DM, Neumann M, Kwong LK, Chou TT, Micsenyi M, Truax A, Bruce J, Grossman M, Trojanowski JQ, Lee VM (2006) Pathological heterogeneity of frontotemporal lobar degeneration with ubiquitin-positive inclusions delineated by ubiquitin immunohistochemistry and novel monoclonal antibodies. *Am J Pathol* 169:1343–1352

46. Seelaar H, Klijnsma KY, de Koning I, van der Lugt A, Chiu WZ, Azmani A, Rozemuller AJ, van Swieten JC (2009) Frequency of ubiquitin and FUS-positive, TDP-43-negative frontotemporal lobar degeneration. *J Neurol* 257:747–753
47. Seelaar H, Schelhaas HJ, Azmani A, Kusters B, Rosso S, Majoor-Krakauer D, de Rijk MC, Rizzu P, ten Brummelhuis M, van Doorn PA, Kamphorst W, Willemsen R, van Swieten JC (2007) TDP-43 pathology in familial frontotemporal dementia and motor neuron disease without Progranulin mutations. *Brain* 130:1375–1385
48. Shi J, Shaw CL, Du PD, Richardson AM, Bailey KL, Julien C, Stopford C, Thompson J, Varma A, Craufurd D, Tian J, Pickering-Brown S, Neary D, Snowden JS, Mann DM (2005) Histopathological changes underlying frontotemporal lobar degeneration with clinicopathological correlation. *Acta Neuropathol* 110:501–512
49. Skibinski G, Parkinson NJ, Brown JM, Chakrabarti L, Lloyd SL, Hummerich H, Nielsen JE, Hodges JR, Spillantini MG, Thüsgaard T, Brandner S, Brun A, Rossor MN, Gade A, Johannsen P, Sorensen SA, Gydesen S, Fisher EM, Collinge J (2005) Mutations in the endosomal ESCRTIII-complex subunit CHMP2B in frontotemporal dementia. *Nat Genet* 37:806–808
50. Snowden JS, Neary D, Mann DM (2002) Frontotemporal dementia. *Br J Psychiatry* 180:140–143
51. Tsuchiya K, Ishizu H, Nakano I, Kita Y, Sawabe M, Haga C, Kuyama K, Nishinaka T, Oyanagi K, Ikeda K, Kuroda S (2001) Distribution of basal ganglia lesions in generalized variant of Pick's disease: a clinicopathological study of four autopsy cases. *Acta Neuropathol* 102:441–448
52. Urwin H, Ghazi-Noori S, Collinge J, Isaacs A (2009) The role of CHMP2B in frontotemporal dementia. *Biochem Soc Trans* 37:208–212
53. van der Zee J, Pirici D, Van Langenhove T, Engelborghs S, Vandenberghe R, Hoffmann M, Pusswald G, Van den Broeck M, Peeters K, Mattheijssens M, Martin JJ, De Deyn PP, Cruts M, Haubenberger D, Kumar-Singh S, Zimprich A, Van Broeckhoven C (2009) Clinical heterogeneity in 3 unrelated families linked to VCP p.Arg159His. *Neurology* 73:626–632
54. van der Zee J, Urwin H, Engelborghs S, Bruylant M, Vandenberghe R, Dermaut B, De PT, Peeters K, Santens P, De Deyn PP, Fisher EM, Collinge J, Isaacs AM, Van BC (2008) CHMP2B C-truncating mutations in frontotemporal lobar degeneration are associated with an aberrant endosomal phenotype in vitro. *Hum Mol Genet* 17:313–322
55. Vance C, Rogelj B, Hortobagyi T, De Vos KJ, Nishimura AL, Sreedharan J, Hu X, Smith B, Ruddy D, Wright P, Ganesalingam J, Williams KL, Tripathi V, Al-Saraj S, Al-Chalabi A, Leigh PN, Blair IP, Nicholson G, de Belleruche J, Gallo JM, Miller CC, Shaw CE (2009) Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323:1208–1211
56. Watts GD, Wymer J, Kovach MJ, Mehta SG, Mumm S, Darvish D, Pestronk A, Whyte MP, Kimonis VE (2004) Inclusion body myopathy associated with Paget disease of bone and frontotemporal dementia is caused by mutant valosin-containing protein. *Nat Genet* 36:377–381
57. Woulfe J, Gray DA, Mackenzie IR (2009) FUS-Immunoreactive Intranuclear Inclusions in Neurodegenerative Disease. *Brain Pathol* (in press)
58. Yokota O, Tsuchiya K, Terada S, Ishizu H, Uchikado H, Ikeda M, Oyanagi K, Nakano I, Murayama S, Kuroda S, Akiyama H (2008) Basophilic inclusion body disease and neuronal intermediate filament inclusion disease: a comparative clinicopathological study. *Acta Neuropathol* 115:561–575